

MD DHMH Assessment Guidance: Exposed Woman

- Living in* or having traveled to area with ongoing Zika virus transmission, OR
- Having condomless sex with a male with travel to an area with ongoing Zika virus transmission

Non-pregnant

Pregnant

No symptoms

Symptoms

- Test and counsel patient to avoid mosquito bites
- If partner exposed, counsel condoms or abstinence

- Do not test
- Counsel patient to avoid mosquito bites
- Pregnancy prevention guidance per CDC table
- No further follow up unless new exposure

- Test and counsel patient to avoid mosquito bites
- Pregnancy prevention guidance per CDC table

Any non-negative result

Negative

Any non-negative result

- Notify provider and LHD
- Counsel patient to avoid mosquito bites
- Confirmatory testing if needed
- Enter case in surveillance databases (NEDSS, ArboNet)

- Counsel patient to avoid mosquito bites
- No further follow up unless new exposure

- Notify provider and direct to CDC Guidance for Follow-up of Pregnant Women
- Notify LHD
- Counsel patient to avoid mosquito bites
- Confirmatory testing if needed
- Enter case in surveillance databases (NEDSS, ArboNet)
- Enter case in pregnancy registry

- Notify MDA and LHD for mosquito control and response:
 - Mosquito surveillance
 - Larvicide application
 - Adulticide application
 - Education
 - Case finding/enhanced surveillance

- Notify MDA and LHD for mosquito control and response:
 - Mosquito surveillance
 - Larvicide application
 - Adulticide application
 - Education
 - Case finding/enhanced surveillance

DRAFT
April 6, 2016
Maryland
Department of
Health and Mental
Hygiene

*** "Living in" could include living in Maryland when local transmission has been identified**

MD DHMH Assessment Guidance: Exposed Man

- Living in* or having traveled to area with ongoing Zika virus transmission, OR
- Having condomless sex with a male with travel to an area with ongoing Zika virus transmission

If exposed man has pregnant partner, test partner regardless of symptoms in the man (see exposed woman algorithm) and counsel patient to use condoms for duration of pregnancy or abstain

Exposed man has no symptoms

- Do not test unless pregnant partner tests positive
- Counsel patient to avoid mosquito bites
- Counsel patient to use condoms for at least 8 weeks after exposure
- No further follow up unless new exposure

Exposed man has symptoms

Test and counsel patient to avoid mosquito bites

Any non-negative result

- Notify provider and LHD
- Counsel patient to avoid mosquito bites
- Reinforce sexual transmission prevention messages - use condoms for at least 6 months after symptoms started
- Confirmatory testing if needed
- Enter case in surveillance databases (NEDSS, ArboNet)
- Notify MDA and LHD for mosquito control and response:
 - Mosquito surveillance
 - Larvicide application
 - Adulticide application
 - Education
 - Case finding/enhanced surveillance

Negative

- Counsel patient to avoid mosquito bites
- Counsel patient to use condoms for at least 8 weeks after exposure
- No further follow up unless new exposure

DRAFT
April 6, 2016
Maryland
Department of
Health and Mental
Hygiene

*** “Living in” could include living in Maryland when local transmission has been identified**